

---

## Palmer LTER: Photoacclimation in a coastal phytoplankton bloom

OSCAR SCHOFIELD, *Institute of Marine and Coastal Sciences, Rutgers University,  
New Brunswick, New Jersey 08903-0231*

MARK M. MOLINE and B.B. PRÉZELIN, *Department of Biology, University of California, Santa Barbara, California 93106*

Phytoplankton blooms in antarctic waters show a strong coherence with oceanic density fields; high biomass is observed when mixing depths are shallow. It has been hypothesized that when phytoplankton are maintained in a stable light environment, cells photoacclimate and overcome the

chronic light-limitation of growth associated with deeply mixing water columns (Mitchell et al. 1991). Phytoplankton photoacclimate by increasing cellular absorption capabilities ( $\bar{a}_{ph}:m^{-1}$ ) and/or the quantum yield for carbon fixation ( $\phi$ :mole carbon fixed per mole photons absorbed) (Kirk 1994,




Figure 1. Contour of chlorophyll-a ( $\text{mg m}^{-3}$ ) from December 1991 through February 1992 at LTER station B (Prézelin et al. 1992). Black circles indicate discrete samples. The mixed layer depth is denoted by a black line.

pp. 290–303); however, documenting photoacclimation in field populations is difficult. As part of the Palmer Long-Term Ecological Research (LTER) project, the temporal dynamics of a phytoplankton bloom was documented near Palmer Station ( $64^{\circ}40'S$   $64^{\circ}03'W$ ) during the austral summer months of 1991–1992 (figure 1). This was an ideal data set for photoacclimation in phytoplankton because we were able to (1) follow the formation of bloom over time and (2) estimate  $\bar{a}_{ph}$  and  $\phi$ .

Water samples were collected from station B within the nearshore sampling grid of the LTER (Prézelin et al. 1992). *In situ* light levels ( $Q_{PAR}$ ) were measured (Biospherical QL-100 scalar irradiance meter), and discrete samples were transported to Palmer Station for analysis. Phytoplankton pigmentation was determined by high-performance liquid chromatography following the protocols described in Prézelin et al. (1992). Phytoplankton absorption was calculated as the summation of the products of *in vivo* absorption coefficients [ $a_i^*(\lambda)$ , figure 2A] and the volume-based concentrations [ $c_i$ , milligrams per cubic meter ( $\text{mg m}^{-3}$ )] of the individual chlorophyll and carotenoid pigments (Bidigare et al. 1990). To calculate the absorption capabilities of the phytoplankton at depth, the reconstructed phytoplankton absorption spectra (figure 2B) must be weighted to the incident spectral irradiance [ $Q(\lambda)$ ].

$$\bar{a}_{ph}' = \frac{\int_{400nm}^{700nm} a_{ph}(\lambda) \cdot Q(\lambda) d\lambda}{Q_{PAR}}$$

Measurements of the incident spectral irradiance were not available; thus, the absorption spectra were weighted to the spectral irradiance measured for oligotrophic coastal waters at the percentage of  $Q_{PAR}$  light depth where the LTER discrete water samples had been collected (Schofield et al. 1993; figure 2C). This results in an upper limit estimate of  $\bar{a}_{ph}$  because  $Q(\lambda)$  in oligotrophic waters will be shifted to the blue-green wavelengths of light where phytoplankton absorption is high. The wavelengths of maximum light transmission in a water column with high concentrations of phytoplankton pre-




Figure 2. Procedure to calculate the spectrally weighted absorption coefficient for phytoplankton. A. The pigment-specific absorption coefficients for the chlorophylls and carotenoids measured by high performance liquid chromatography. B. An example of a reconstructed phytoplankton absorption spectra. C. The spectral irradiance profile used to weight the reconstructed absorption spectra to the incident light quality.

sent (as during the bloom) will be shifted to the green-orange wavelengths of light where cellular absorption is lower (figure 2B). Chlorophyll-specific  $\bar{a}_{ph}$  showed no correlation with the phytoplankton bloom (figure 3A), a finding that provides circumstantial evidence that modifications in cellular absorption capabilities do not constitute the primary mode of photoacclimation for these phytoplankton; however, it should be emphasized that if cells increased all pigments in equal molar ratios then the chlorophyll-specific  $\bar{a}_{ph}$  would remain relatively constant. We are currently assessing this possibility by studying the temporal variability in the accessory pigment-chlorophyll-*a* ratios.

Values of  $\phi$  were estimated from productivity rates calculated from photosynthesis-irradiance curves by using procedures described by Schofield et al. (1993). Values of  $\phi$  increased prior to the phytoplankton bloom (figure 3B) and values approached theoretical maximum values as the mixed layer depth shallowed. This suggests that the regulation of  $\phi$  was a primary mode of photoacclimation. Later in the season, as the mixed layer depth shallowed, the values of  $\phi$  increased. These data suggest that photoacclimation in these phytoplankton was constrained by water column mixing, which is consistent

with the hypothesis that shallow mixing depths provide a relatively stable light environment that allows algae to photoacclimate on the timescale of 1–2 days.

This work was supported by National Science Foundation grants OPP 90-11927 and OCE 93-01322 awarded to Barbara B. Prézelin. This is Palmer LTER contribution number 59.

### References

- Bidigare, R.R., M.E. Ondrusek, M.E. Morow, and D.A. Kiefer. 1990. *In vivo* absorption properties of algal pigments. *Proceedings SPIE Ocean Optics X*, 1302, 290–297.
- Kirk, J.T.O. 1994. *Light and photosynthesis in aquatic ecosystems* (2nd ed.). Cambridge: Cambridge University Press.
- Mitchell, B.G., E.A. Brody, O. Holm-Hansen, C. McClain, and J. Bishop. 1991. Light-limitation of phytoplankton biomass and macronutrient utilization in the southern ocean. *Limnology and Oceanography*, 36(8), 1662–1677.
- Prézelin, B.B., M. Moline, K. Sydell, and K. Scheppe. 1992. Palmer LTER: Temporal variability in HPLC pigmentation and inorganic nutrient distribution in surface waters adjacent to Palmer Station, December 1991–February 1992. *Antarctic Journal of the U.S.*, 28(5), 245–248.
- Schofield, O., B.B. Prézelin, R.R. Bidigare, and R.C. Smith. 1993. *In situ* photosynthetic quantum yield. Correspondence to hydrographic and optical variability within the southern California Bight. *Marine Ecology Progress Series*, 93, 25–37.




Figure 3. Contours of (A) the chlorophyll-specific spectrally weighted absorption coefficient ( $\text{m}^2 \text{mg chl a}^{-1}$ ) and (B) the operational quantum yield for carbon fixation.